



Activity #2

# Impacts of Aquarium Fish Collecting on Coral Reefs

## ● ● ● In Advance *Student Assignment*

- As homework, assign the Student Pages “Impacts of Aquarium Fish Collectors on Coral Reef Fishes in Kona, Hawai‘i” (pp. 34-41) and “Questions About the Reading” (pp. 42-45).

## ● ● ● Class Period One *Discussing Study Designs*

### Materials & Setup

---

*For each student*

- Student Page “Impacts of Aquarium Fish Collectors on Coral Reef Fishes in Kona, Hawai‘i” (pp. 34-41)
- Student Page “Questions About the Reading” (pp. 42-45)

### Instructions

---

- 1) Beginning with student questions and responses to the homework assignment, hold a class discussion about the study, its design, conclusions, and implications for managing coral reef fisheries.
- 2) For advanced classes, move from the general discussion to a more detailed discussion of experimental design. For guidance and background, use “Impacts of Aquarium Collectors on Coral Reef Fishes in Kona, Hawai‘i” (complete original report in appendix) and “Notes for Class Discussion on ‘Impacts of Aquarium Collectors on Coral Reef Fishes’” (pp. 28-29).

The class discussion should get into the details of experimental design at an appropriate depth for the level of students. Students should be prepared for this discussion based on their reading and answering the homework questions.
- 3) Wrap up the discussion by focusing on the final homework question, how students would go about learning whether aquarium collecting is a current problem on Maui and what they would do to prevent it from becoming a problem in the future.

### Journal Ideas

---

- If you were in charge of regulating aquarium fish collecting, what would you do based on reading the Tissot and Hallacher study?
- Do you think that people should be allowed to collect native Hawaiian fish for the aquarium trade? Why or why not?

### Assessment Tools

---

- Student Page “Questions About the Reading” (teacher version, pp. 30-33)
- Participation in class discussion
- Journal entries



### *Teacher Background*

## Notes for Class Discussion on “Impacts of Aquarium Collectors on Coral Reef Fishes”

To make the main points of this paper easier for students to grasp, several parts of the study and discussion were left out of the simplified student version of the paper. Depending upon the level of student and the nature of the course you are teaching, you will want to go into details of the study that will help students understand experimental design.

Below are the main elements of the paper, along with page references to the full version of the paper (see appendix). First, make sure that students understand the results of the study as presented in the student version. Then, with the time remaining, go into the elements of experimental design, emphasizing the first three sections of the paper.

#### 1) Introduction/statement of the problem (pp. 4-5)

The context for this study is the growing global and Hawai‘i-based trade in marine aquarium fishes collected from the wild. The authors identified a lack of conclusive studies documenting the magnitude of impacts on natural populations.

#### 2) Scope/purpose of the study (p. 4)

This study had two main purposes. Only the first is covered in the student version of the paper:

- a) Obtain quantitative estimates of the impact of aquarium collectors on reef fishes;
- b) Evaluate evidence for destructive harvesting methods and changes in the reef community associated with reductions in herbivory (predation on plants).

#### 3) Methods (pp. 5-7)

- a) The authors explain their experimental design (a paired control-impact design) and its major assumptions. The student version of the paper deals very little with the assumptions of the design and how they were tested/addressed. The main assumptions and how they were tested and addressed in the study and report are:

#### Assumption

Prior to the onset of aquarium harvesting, there was no difference in abundance of aquarium fishes between the control and impact sites. (The study was begun after the impacts had begun.)

#### How addressed, tested

- Paired control and impact sites were geographically close together (p. 10).
- The survey was conducted on corals, macroalgae, and the general substratum of each transect. Analysis showed remarkable similarity between the paired control-impact sites (pp. 8, 10).
- The survey included ecologically similar species not targeted by aquarium collectors. A prediction of this assumption is that non-collected species should not differ between control and impact sites. This prediction was supported by study data (pp. 6, 8-10).



## Assumption

All differences between the control and impact sites are due to aquarium fish collecting and not other factors, such as fishing.

## How addressed, tested

- Impact sites were largely inaccessible from shore, minimizing shore-based recreational fishing (p. 10).
- There was no significant control-impact variation in abundance of a nontarget species (not collected for aquariums) subject to commercial and recreational fishing (p. 10).
- Introduced piscivorous (fish-eating) fishes that may cause significant mortality among aquarium species were rare at the study sites (pp. 10-11).

- b) The authors also describe their research sites, methods, personnel, and timing of the surveys, as well as steps they took to minimize bias (pp. 6-7). Much of this detail has been left out of the student version of the paper.
- c) Methods of analysis are described (p. 7). Only the basic calculation of percent change in abundance is described in the student version of the paper. The analysis for statistical significance is not described in the student version, nor were the factors included (impact, location, and impact-location).
- 4) Results (pp. 6-9)  
In the student version of the paper, this section is condensed and combined with the discussion section, primarily because so many of the results had to do with analyses not covered in the student paper.
- 5) Discussion (pp. 9-13)  
In this paper, the discussion of results has four sections:  
-Evaluation of assumptions (see table above)  
-Magnitude of impacts (eliminated from the student version)  
-Indirect effects (eliminated from the student version)  
-Implications for fishery management
- 6) Acknowledgments (p. 13)
- 7) References  
References in the student version have been modified based on editing the original.
- 8) Tables and figures  
Only some of the tables from the original paper are in the student version.



*Teacher Version*

## Questions About the Reading

Answer the following questions. Attach additional sheets if necessary.

- 1) What was the purpose of the study?

To examine the effects of aquarium collecting on reef fish populations in Hawai'i

- 2) Why did the researchers choose to survey both fish species that are commonly collected for the aquarium trade and those that are not?

Each nontargeted fish species was chosen because its habitat and food type are similar to one or more targeted species. These nontarget species provided a base of comparison that helped researchers determine whether the changes in population were related to aquarium collecting.

- 3) Explain what the “impact” sites were and what the “control” sites were and why they are important in this study.

The impact sites were areas with high levels of aquarium collecting. The control sites were adjacent to impact sites, in areas where aquarium collecting is prohibited. Researchers could estimate the magnitude of impact by comparing population density and changes in population density at sites where aquarium collecting occurs and where it does not.



- 4) The researchers were interested in determining how much difference there was between fish abundance at the control and impact sites. They determined the mean density of fish at each of the sites. Then they calculated a percent change in fish abundance for each species, and for each species at each of the two study sites.

A negative percent change indicates fewer fish at the impact relative to the control site, while a positive value indicates the opposite pattern.

Species	Mean overall percent change
<b>Aquarium fishes</b>	
Achilles tang ( <i>Acanthurus achilles</i> )	-57.1
Potter's angelfish ( <i>Centropyge potteri</i> )	-46.1
Multi-band butterflyfish ( <i>Chaetodon multicinctus</i> )	-38.2
Ornate butterflyfish ( <i>Chaetodon ornatissimus</i> )	-39.5
Four-spot butterflyfish ( <i>Chaetodon quadrimaculatus</i> )	-41.6
Goldring surgeonfish ( <i>Ctenochaetus strigosus</i> )	-14.7
Longnose butterflyfish ( <i>Forcipiger</i> spp.)	-54.2
Orangespine unicornfish ( <i>Naso lituratus</i> )	31.2
Moorish idol ( <i>Zanclus cornutus</i> )	-46.5
Yellow tang ( <i>Zebrasoma flavescens</i> )	-47.3
<b>Non-Aquarium Species</b>	
Brown surgeonfish ( <i>Acanthurus nigrofuscus</i> )	27.3
Blueline surgeonfish ( <i>Acanthurus nigroris</i> )	67.2
Convict surgeonfish or tang ( <i>Acanthurus triostegus</i> )	-4.3
Oval butterflyfish ( <i>Chaetodon lunulatus</i> )	-70.0
Arc-eye hawkfish ( <i>Paracirrhites arcatus</i> )	-36.4
Blackside hawkfish ( <i>Paracirrhites forsteri</i> )	58.4
Blue-eye damsel ( <i>Plectroglyphidodon johnstonianus</i> )	-31.3
Pacific gregory ( <i>Stegastes fasciolatus</i> )	326.0
Saddle wrasse ( <i>Thalassoma duperrey</i> )	17.4

- 4a) Which three species show the greatest difference between the number of individuals at control sites and impact sites? For each species, identify whether this difference indicates that there are fewer individuals at the control sites or the impact sites.

Pacific gregory (*Stegastes fasciolatus*) — fewer at control sites

Oval butterflyfish (*Chaetodon lunulatus*) — fewer at impact sites

Blueline surgeonfish (*Acanthurus nigroris*) — fewer at control sites



Activity #2  
Marine Unit 5

---

- 4b) Which four species show the greatest negative mean percent change—indicating fewer individuals at the impact sites relative to the control sites? Discuss the possible significance of these results based on whether these species are collected for the aquarium trade or not.

Oval butterflyfish (*Chaetodon lunulatus*)  
Achilles tang (*Acanthurus achilles*)  
Longnose butterflyfish (*Forcipiger spp.*)  
Yellow tang (*Zebrasoma flavescens*)

Three of the four species are targeted by aquarium collectors. This probably indicates that aquarium collection, which occurs at the impact sites and not the control sites, decreases fish populations.

- 5) What patterns do you notice when you compare the aquarium species with the non-aquarium species, looking at whether the percent change is negative or positive? What do these patterns suggest about the impact of aquarium collecting?

All but one of the aquarium species show a negative percent change, while the results are mixed among non-aquarium species (three negative and four positive). This result suggests that aquarium collecting decreases populations of target species and that other factors might also have come into play (because of the mixed results among the non-aquarium species).

- 6) The experimental design that the researchers selected for this study makes two major assumptions:
- The study began after aquarium fish collecting had already started in the impact areas. Therefore, the design assumes that the natural abundance of aquarium fishes at the control and impact sites were similar prior to the onset of aquarium collection.
  - The design assumes that all differences between the paired control and impact sites were due to aquarium fish collecting and not other factors, such as fishing.

Choose one of these assumptions and think of a way that the researchers could — or did — build into the study a way to test whether the assumption seems valid.

There are many correct responses to this question, which should be evaluated based on student reasoning, as well as references back to the study:



a) One way the researchers controlled for the first assumption was to select control and impact sites that are close together, to lessen the likelihood of spatial variation. They also used a combination of nontarget species that were ecologically similar to target species, and other species that were indicators of particular habitats, as indicators of the ecological similarity of the control and impact sites.

Other ways of assessing the correctness of this assumption include comparing control and impact sites for factors such as species diversity and richness, and comparing the habitats by looking at coral and algae abundance and diversity, and non-living substratum composition.

b) One way to control for the second assumption is to select sites that are largely inaccessible from shore, to minimize the impact of shore-based recreational fishing. (The authors did this in the aquarium collecting study, but that may not be clear from the student background reading.)

Other ways of testing the assumption include looking at density variation in target and nontarget species that are subject to commercial and recreational fishing. If the density of nontarget species that are fished for does not vary between control and impact sites, this suggests that fishing impacts are not significant.

Another factor that could differentially affect reef fish populations could be the presence of predator fishes. Including predator fishes in the surveys would help determine whether this factor does influence reef fish populations.

- 7) Some people say that aquarium collecting is not a problem on Maui, while others believe that it is a problem in some areas or could quickly become one. Write one paragraph about what you would do to find out whether aquarium collecting is a threat to Maui reef animals. Write another paragraph about what you think should be done, if anything, to protect Maui reef fish populations from the impacts of collecting.

Well-reasoned responses are acceptable.



# Impacts of Aquarium Collectors on Coral Reef Fishes in Kona, Hawai'i

*This paper is condensed with authors' permission from a September 1999 report of the same name prepared for the State of Hawai'i Division of Aquatic Resources by Brian N. Tissot, Ph.D. (Washington State University — Vancouver) and Leon E. Hallacher, Ph.D. (University of Hawai'i — Hilo). This distillation of their paper focuses on the first goal of their study: to obtain "quantitative" or number-based estimates of the impact of aquarium collectors on reef fishes. It does not cover methods, data, or conclusions associated with the second goal of the study: to evaluate evidence for destructive fish harvesting methods and changes in the reef community associated with reductions in herbivory.*

## Introduction

Each year, some 350 million ornamental aquarium fish worth \$963 million are sold around the world (Young, 1997). Although marine fishes account for only ten to 20 percent of the total, the harvest level for marine species grew rapidly in the 1980s (Andrews, 1990). Over 99 percent of marine fishes sold in the aquarium trade are taken from the wild, unlike their freshwater counterparts, most of which are cultivated (Young, 1997). Almost all marine ornamentals are of tropical origin and many are harvested from coral reefs. Because aquarium fish collectors focus heavily on a few species and often capture large quantities of individuals of high value, the potential for overfishing is high (Wood, 1985).

Many studies have discussed the potential effects of the aquarium trade on marine fishes in Australia (Whitehead et al., 1986), Hawai'i (Taylor, 1978; Walsh, 1978; Randall, 1987), Indonesia (Wood, 1985), the Philippines (Albaladejo and Corpuz, 1981), Puerto Rico (Sadovy, 1992), and Sri Lanka (Edwards and Shepherd, 1992). But there are no conclusive studies documenting the magnitude of impacts on fish populations, despite repeated calls for such studies to help sustain the aquarium trade industry over the long term (Walsh, 1978; Wood, 1985; Young, 1997).

Most of the marine ornamentals originating from the U.S. are taken from Hawai'i waters. Hawai'i is known for its high-quality fishes and rare endemic fishes of high value (Wood, 1985). As early as the 1970s, concerns over the effects of aquarium collecting on reef fish populations in Hawai'i were being raised. (Taylor, 1978; Walsh, 1978). Aquarium fish collectors and recreational dive tour operators came into conflict over apparent declines in nearshore reef fishes (Taylor, 1978). This conflict continues up to the present (Grigg, 1997; Young, 1997; Clark and Gulko, 1999). Early concerns prompted the Hawai'i Division of Fish and Game [now the Division of Aquatic Resources] to require monthly collection reports of all permit holders starting in 1973 (Katekaru, 1978). These reports have been the primary basis for managing the aquarium industry in Hawai'i since then (Miyasaka, 1994, 1997).

Data from collection reports suggest that the size and value of the Hawai'i aquarium fish industry is growing. In 1973, 90,000 fishes with a total value of \$50,000 were reported (Katekaru, 1978). In 1995, the annual harvest had risen to 422,823 fishes with a total value of \$844,843 (Miyasaka, 1997).

Although a total of 103 fish species were collected statewide in 1995 (Division of Aquatic Resources [DAR], unpublished data), over 90





percent of the harvest is focused on 11 species. The yellow tang (*Zebrasoma flavescens*) accounted for 52 percent of the total harvest in 1995 (DAR, unpublished data; Miyasaka, 1997). Given the increasing rate of harvest focused on a small number of species, the potential for “overexploitation” is high, meaning the fishes are taken at such a rate that they cannot maintain their populations over the long term.

## Materials and Methods

We used a “paired control-impact design” to estimate the impact of aquarium collectors on the “abundance” or relative numbers of reef fish in an area. The magnitude of the impact was estimated by comparing the difference between fish abundance at “impact” sites, where aquarium fish collecting was known to occur, relative to nearby “control” sites where collecting was prohibited.

We established four study sites that served as two control-impact pairs for the study (Figure 1). Impact sites were selected in areas where high levels of aquarium fish collecting was occurring (personal communications). Control sites were located in areas adjacent to impact sites, where aquarium fish collecting was prohibited.

The first pair of study sites were located at Honokōhau and Papawai on the island of Hawai‘i. Papawai is a Fishery Management Area (FMA) where the collecting of aquarium fishes is prohibited (DLNR, 1996). It served as a control site. Honokōhau was frequented by aquarium collectors and served as an impact site. These paired sites will hereafter be referred to as the “Honokōhau” study area. The second pair of study sites were located at Red Hill North and Red Hill South. Red Hill South is a FMA

where the collecting of aquarium fishes is prohibited (DLNR, 1996), and which served as a control site. Red Hill North was frequented by aquarium collectors and served as an impact site. These paired sites will hereafter be referred to as the “Red Hill” study area.

At each study site four permanent 50-meter “transects” or lines were established at ten to 15 meter depths by installing stainless steel eyebolts at the beginning and end points of each. The abundances of fishes was estimated using a visual strip-transect search method (Sale and Douglas, 1981). In this method, a pair of divers swam side-by-side down either side of the transect line and count all fish seen within a corridor three meters wide and extending to the surface.

Surveys began at Honokōhau in March, and at Red Hill in September, of 1997 and ended at both sites in December 1998. All sites were sampled at




Figure 1. Map of study sites located on the island of Hawai‘i



intervals of two to five months for a total of eight surveys at Honokōhau and five at Red Hill.

During each survey we estimated the abundance of 21 fish species (Table 1). Eleven aquarium fish species were selected based on reported high levels of collection. In addition, we also surveyed ten fish species not targeted by aquarium collectors. These species were selected to serve as indicators of specific habitats and food types and provide data to support the study's assumptions.

The divers used in this study were undergraduates who had completed a rigorous coral reef-monitoring course and were trained in species identification and survey techniques (Russell, 1997; Hallacher and Tissot, 1999). In order to minimize observer bias, the same diver-pairs were used at each control-impact study site during each survey. Divers did, however, vary among surveys. To minimize variation, all surveys were conducted in the middle of the day (generally from 9:00 a.m. to 3:00 p.m.) and both control and impact sites were surveyed either on the same day or on consecutive days.

Percent change in fish abundance was calculated as the difference between both control sites and both impact sites using the formula:

$$\text{Percent change} = [(D_{\text{impact}} - D_{\text{control}}) / D_{\text{control}}] \times 100$$

Where D = density expressed as number of individuals per 100 square meters. Thus, a negative percent change indicates fewer fish at the impact relative to the control sites, while a positive value indicates the opposite pattern.

## Results and Discussion

Of the 21 species surveyed, two species (Raccoon butterflyfish, *Chaetodon lunula*, and Teardrop butterflyfish, *C. unimaculatus*) were too rare for analysis with one individual of each species observed during the entire study. These species were excluded from any further analysis.

Overall, there were numerous "significant" differences (which are unlikely to occur based on chance alone) in the abundance of aquarium

fishes between control and impact sites but few differences in the abundance of non-aquarium species (Table 2).

The results of this study indicate that eight of the ten fishes targeted by aquarium collectors were significantly reduced in abundance in areas subjected to harvesting, relative to managed areas where collecting was prohibited. The magnitude of these declines ranged from 57 percent in Achilles tang (*Acanthurus achilles*) to 38 percent in Multi-band butterflyfish (*Chaetodon multicinctus*). In contrast, only one of the nine "nontarget" species not typically collected for aquariums varied significantly between these areas, suggesting that aquarium collectors are having significant impacts on the abundance of targeted fishes in near-shore areas on the Kona coast of Hawai'i.

## Evaluation of Assumptions

Part of the design of this study was to use a combination of nontargeted species that were ecologically similar to target species (those that are collected for the aquarium trade). This is one way to infer whether observed differences are due to the impact of aquarium collectors or due to other differences between the control and impact sites.

Overall, aquarium fishes exhibited significant differences between control and impact sites, while nontarget species did not. Table 3 details some of these comparisons.

The one exception to this pattern was the Arc-eye hawkfish (*Paracirrhites arcatus*), the only nontarget species that was significantly less abundant in impact relative to control areas. This species lives in close association with corals, primarily *Pocillopora meandrina*, which although rare at all study sites, was less abundant at impact relative to control sites.

## Implications for Fishery Management

This study indicates that aquarium collectors are having significant impacts on eight of the ten  
(Continued on p. 40)



Table 1. List of fishes monitored during the study

Information on diet and trophic level is based on Randall (1985, 1996).

SPECIES	TROPHIC LEVEL	DIET
<i>Aquarium fishes</i>		
Achilles tang ( <i>Acanthurus achilles</i> )	Herbivore	Filamentous algae
Potter's angelfish ( <i>Centropyge potteri</i> )*	Herbivore	Filamentous algae and detritus
Raccoon butterflyfish ( <i>Chaetodon lunula</i> ) <sup>o</sup>	Carnivore	Small invertebrates
Multi-band butterflyfish ( <i>Chaetodon multicinctus</i> )*	Corallivore	Coral polyps
Ornate butterflyfish ( <i>Chaetodon ornatissimus</i> )	Corallivore	Coral polyps
Four-spot butterflyfish ( <i>Chaetodon quadrimaculatus</i> )	Corallivore	Coral polyps
Goldring surgeonfish ( <i>Ctenochaetus strigosus</i> )	Detritivore	Detritus
Longnose butterflyfish ( <i>Forcipiger</i> spp.)*	Carnivore	Small invertebrates
Orangespine unicornfish ( <i>Naso lituratus</i> )	Herbivore	Macroalgae
Moorish idol ( <i>Zanclus cornutus</i> )	Omnivore	Sponges and algae
Yellow tang ( <i>Zebrasoma flavescens</i> )	Herbivore	Filamentous algae
<i>Nonaquarium fishes</i>		
Brown surgeonfish ( <i>Acanthurus nigrofuscus</i> )	Herbivore	Filamentous algae
Blueline surgeonfish ( <i>Acanthurus nigroris</i> ) *	Herbivore	Filamentous algae
Convict tang ( <i>Acanthurus triostegus</i> )	Herbivore	Filamentous algae
Teardrop butterflyfish ( <i>Chaetodon unimaculatus</i> ) <sup>o</sup>	Corallivore	Coral polyps
Oval butterflyfish ( <i>Chaetodon lunulatus</i> )	Corallivore	Coral polyps
Arc-eye hawkfish ( <i>Paracirrhites arcatus</i> )	Carnivore	Invertebrates and fishes
Blackside hawkfish ( <i>Paracirrhites forsteri</i> )	Carnivore	Invertebrates and fishes
Blue-eye damsel ( <i>Plectroglyphidodon johnstonianus</i> )	Corallivore	Coral polyps
Pacific gregory ( <i>Stegastes fasciolatus</i> )	Herbivore	Filamentous algae and detritus
Saddle wrasse ( <i>Thalassoma duperrey</i> )*	Carnivore	Invertebrates

\* endemic to Hawai'i      <sup>o</sup> too rare to be included in the analysis

\* two species of longnose butterflyfish were included in this category



Table 2. Mean density of aquarium and nonaquarium fishes at control and impact study sites pooled for the entire study

	Density (no. / 100 m <sup>2</sup> )			
	Honokōhau		Red Hill	
	Impact	Control	Impact	Control
<b>Aquarium fishes</b>				
<i>Acanthurus achilles</i>	0.23	0.69	0.40	0.92
<i>Centropyge potteri</i>	1.48	2.50	0.25	0.85
<i>Chaetodon multicinctus</i>	2.98	4.95	3.43	5.72
<i>Chaetodon ornatissimus</i>	0.25	0.59	0.57	1.37
<i>Chaetodon quadrimaculatus</i>	0.01	0.15	0.17	0.38
<i>Ctenochaetus strigosus</i>	24.10	35.60	32.10	28.70
<i>Forcipiger spp.</i>	1.27	3.24	0.75	1.33
<i>Naso lituratus</i>	1.58	1.25	0.92	1.72
<i>Zanclus cornutus</i>	0.34	0.89	0.28	0.65
<i>Zebrasoma flavescens</i>	9.72	19.80	14.30	24.40
<b>Overall Density</b>	42.00	69.7	53.20	66.00
<b>Nonaquarium fishes</b>				
<i>Acanthurus nigrofuscus</i>	12.10	11.30	23.90	17.60
<i>Acanthurus nigroris</i>	1.24	2.60	3.42	1.68
<i>Acanthurus triostegus</i>	0.16	0.32	0.17	0.13
<i>Chaetodon lunulatus</i>	0.26	0.11	0.00	0.00
<i>Paracirrhites arcatus</i>	1.28	1.56	0.87	3.68
<i>Paracirrhites forsteri</i>	0.42	0.17	0.15	0.60
<i>Plectroglyphidodon johnstonianus</i>	1.82	2.11	0.97	1.93
<i>Stegastes fasciolatus</i>	1.29	0.73	0.15	0.10
<i>Thalassoma duperrey</i>	3.91	3.22	3.30	3.65
<b>Overall Density</b>	22.50	22.20	32.90	29.40



Table 3: Comparisons in change in abundance among similar target and nontarget species

Species	Similar characteristics	Change in abundance
<u>Nontarget</u> Brown surgeonfish ( <i>Acanthurus nigrofuscus</i> )	Generalized herbivores that feed on filamentous algae, occupy the same depth ranges and habitats, and exhibit similar patterns of spawning and larval recruitment (Randall, 1985; Walsh, 1987; Lobel, 1989)	No significant variation between impact and control sites
<u>Target</u> Yellow tang ( <i>Zebrasoma flavescens</i> )		Forty-seven percent less abundant at impact sites than at control sites
<u>Nontarget</u> Oval butterflyfish ( <i>Chaetodon lunulatus</i> ) Blue-eye damselfish ( <i>Plectroglyphidodon johnstonianus</i> )	Feed on coral or live in close association with coral	No significant variation between impact and control sites
<u>Target</u> Multi-band butterflyfish ( <i>Chaetodon multicinctus</i> ) Ornate butterflyfish ( <i>C. ornatissimus</i> ) Four-spot butterflyfish ( <i>C. quadrimaculatus</i> )		Significantly lower abundances at impact sites
<u>Nontarget</u> Blueline surgeonfish ( <i>Acanthurus nigroris</i> ) Convict surgeonfish ( <i>A. triostegus</i> ) Blackside hawkfish ( <i>Paracirrhites forsteri</i> ) Pacific gregory ( <i>Stegastes fasciolatus</i> ) Saddle wrasse ( <i>Thalassoma duperrey</i> )	Generalized diets and distributions across the reef	No significant variation between impact and control sites
<u>Target</u> Achilles tang ( <i>Acanthurus achilles</i> ) Potter's angelfish ( <i>Centropyge potteri</i> ) Moorish idol ( <i>Zanclus cornutus</i> )		Significantly lower abundances at impact sites



species examined. However, more specific information about location, catch and effort is essential to verify the results of this study. The current system of catch reporting in Hawai‘i is limited to monthly collecting reports, with the 235-kilometer (146-mile) coastline of west Hawai‘i divided into three large sections (Miyasaka, 1997). These reports are not compared to actual catches, so there is no quality assurance that the reports are accurate. Analysis of the current catch reports indicates that significant numbers of reports are not filed (DAR, personal communication). Routine monitoring of the collector’s catch report should be instituted to provide some level of quality assurance about the reported catch data.

The magnitude and extent of the impacts documented in this study clearly point to an increased need for management of these species in Hawai‘i. Responding to continued strong public outcry over the aquarium collecting issue, the Hawai‘i state legislature passed a bill in 1998 which focused on improving management of reef resources. The law established the West Hawai‘i Regional Fishery Management Area. It also set aside a minimum of 30 percent of the west Hawai‘i coastline as Fish Replenishment Areas (FRAs), protected areas where aquarium fish collecting is prohibited. Based largely on input from the West Hawai‘i Fishery Council, a community-based group of individuals, a network of nine FRAs has been proposed as a plan to manage the aquarium industry. Our current efforts are focused on monitoring these areas in order to evaluate the effectiveness of the proposed reserve network as a fishery management tool. Through monitoring of changes in abundance in the reserves relative to existing protected and impact areas (including the Honokōhau and Red Hill study sites), we will be able to test predictions derived from the results of this study.

## Acknowledgments

We thank Brent Carmen, John Kahiapo and John Coney for logistical support during this project.

The actual fish counts were conducted by Rob Ames, Anne Creason, Brent Larsen, Bryan Doo, John Holland, Greg Polloi, Jill Sommer, Monica Gregorita, Michelle Pico, and Rachelle Spears. Jonathan Hultquist analyzed all of the photoquadrat images. Robert Nishimoto, Peter Hendricks, Sara Peck, and David Tarnas assisted in planning and design. The research was supported from a grant from the Hawai‘i Division of Aquatic Resources.

## References

- Albaladejo, V. D., and V. T. Corpuz, “A Market Study of the Aquarium Fish Industry of the Philippines: An Assessment of the Growth and the Mechanics of the Trade,” *Proceedings of the 4th International Coral Reef Symposium*, Vol. 1, 1981, pp. 75-81.
- Andrews, C., “The Ornamental Fish Trade and Fish Conservation,” *Journal of Fish Biology*, Vol. 37, Supplement A, 1990, pp. 53-59.
- Clark, A. M., and D. Gulko, *Hawai‘i’s State of the Reefs Report, 1998*, Department of Land and Natural Resources, Honolulu, 1999.
- DLNR, *Hawai‘i Fishing Regulations*, Division of Aquatic Resources, Honolulu, 1996.
- Edwards, A. J., and A. D. Shepherd, “Environmental Implications of Aquarium-Fish Collection in the Maldives, with Proposals for Regulation,” *Environmental Conservation*, Vol. 19, 1992, pp. 61-72.
- Grigg, R. W., “Hawaii’s Coral Reefs: Status and Health in 1997, the International Year of the Reef,” in R.W. Grigg and C. Birkeland (eds.), *Status of Coral Reefs in the Pacific*, University of Hawai‘i Sea Grant College Program, Honolulu, 1997, pp. 59-72.



Hallacher, L. E., and B. N. Tissot, "QUEST: Quantitative Underwater Ecological Survey Techniques: A Coral Reef Monitoring Workshop," *Proceedings of the Hawai'i Coral Reef Monitoring Workshop*, Department of Land and Natural Resources, Honolulu, 1999.

Katekaru, A., "Regulations of Tropical Fish Collecting," *Working Papers No. 34*, Hawai'i Sea Grant College Program, Honolulu, 1978, pp. 35-42.

Miyasaka, A., *Status Report, Aquarium Fish Collections, Fiscal Year 1993-1994*, Division of Aquatic Resources, Honolulu, 1994.

Miyasaka, A., *Status Report, Aquarium fish Collections, Fiscal Year 1994-95*, Division of Aquatic Resources, Honolulu, June 1997.

Randall, J. E., "Collecting Reef Fishes for Aquaria," in B. Salvat (ed.), *Human Impacts on Coral Reefs: Facts and Recommendations*, Antenne Museum EPHE, French Polynesia, 1987, pp. 29-39.

Russell, S. H., "Coral Reef Conservation and Management: Providing Scientific Training to the Community," *Proceedings of the 8th International Coral Reef Symposium*, Vol. 2, 1997, pp. 2087-2088.

Sadovy, S., "A Preliminary Assessment of the Marine Aquarium Export Trade in Puerto Rico," *Proceedings of the 7th International Coral Reef Symposium*, Vol. 2, 1992, pp. 1014-1022.

Sale, P. F., and W. A. Douglas, "Precision and Accuracy of Visual Census Techniques for Fish Assemblages on Coral Reef Patch Reefs," *Env. Biol. Fish.*, Vol. 6, 1981, pp. 333-339.

Taylor, L. R., "Tropical Reef Fish Management: Issues and Opinions," *Working Papers No. 34*, Hawai'i Sea Grant College Program, 1978, pp. 3-5.

Walsh, W. J., "Aquarium Fish Collecting: Promise or Problem?" *Working Papers No. 34*, Hawai'i Sea Grant College Program, 1978, pp. 8-12.

Whitehead, M. J., E. Gilmore, P. Eager, P. McMinnity, W. Craik, and P. Macleod, *Aquarium Fishes and Their Collection in the Great Barrier Reef Region*, Great Barrier Reef Marine Park Authority Technical Memorandum GBRMPA-TM-13, 1986.

Wood, E., *Exploitation of Coral Reef Fishes for the Aquarium Trade*, Report to Marine Conservation Society, United Kingdom, 1985.

Young, L. G. L., "Sustainability Issues in the Trade for Wild and Cultured Aquarium Species," in *Marketing and Shipping Live Aquatic Products, '96, Seattle, Washington, 13-15 October, 1996* (conference proceedings), Northeastern Regional Agricultural Engineering Service Cooperative Extension, Ithaca, New York, 1997, pp. 145-151.







- 4) The researchers wanted to determine how much difference there was between fish abundance at the control and impact sites. They determined the mean density of fish at each of the sites. Then they calculated a percent change in abundance for each species.

A negative percent change indicates fewer fish at the impact relative to the control site, while a positive value indicates the opposite pattern.

Species	Mean percent change for both study sites
---------	---

**Aquarium fishes**

Achilles tang ( <i>Acanthurus achilles</i> )	-57.1
Potter's angelfish ( <i>Centropyge potteri</i> )	-46.1
Multi-band butterflyfish ( <i>Chaetodon multicinctus</i> )	-38.2
Ornate butterflyfish ( <i>Chaetodon ornatissimus</i> )	39.5
Four-spot butterflyfish ( <i>Chaetodon quadrimaculatus</i> )	-41.6
Goldring surgeonfish ( <i>Ctenochaetus strigosus</i> )	14.7
Longnose butterflyfish ( <i>Forcipiger</i> spp.)	-54.2
Orangestripe unicornfish ( <i>Naso lituratus</i> )	31.2
Moorish idol ( <i>Zanclus cornutus</i> )	-46.5
Yellow tang ( <i>Zebrasoma flavescens</i> )	-47.3

**Nonaquarium Species**

Brown surgeonfish ( <i>Acanthurus nigrofuscus</i> )	27.3
Blueline surgeonfish ( <i>Acanthurus nigroris</i> )	67.2
Convict tang ( <i>Acanthurus triostegus</i> )	-4.3
Oval butterflyfish ( <i>Chaetodon lunulatus</i> )	-70.0
Arc-eye hawkfish ( <i>Paracirrhites arcatus</i> )	-36.4
Blackside hawkfish ( <i>Paracirrhites forsteri</i> )	58.4
Blue-eye damsel ( <i>Plectroglyphidodon johnstonianus</i> )	-31.3
Pacific gregory ( <i>Stegastes fasciolatus</i> )	326.0
Saddle wrasse ( <i>Thalassoma duperrey</i> )	17.4

- 4a) Which three species show the greatest difference between the number of individuals at control sites and impact sites? For each species, identify whether this difference indicates that there are fewer individuals at the control sites or the impact sites.



- 4b) Which four species show the greatest negative mean percent change—indicating fewer individuals at the impact sites relative to the control sites? Discuss the possible significance of these results based on whether these species are collected for the aquarium trade or not.
- 5) What patterns do you notice when you compare the aquarium species with the non-aquarium species, looking at whether the percent change is negative or positive? What do these patterns suggest about the impact of aquarium fish collecting?
- 6) The experimental design that the researchers selected for this study makes two major assumptions:
- The study began after aquarium fish collecting had already started in the impact areas. Therefore, the design assumes that the natural abundance of aquarium fishes at the control and impact sites were similar prior to the onset of aquarium collection.
  - The design assumes that all differences between the paired control and impact sites were due to aquarium fish collecting and not other factors, such as sport fishing or pollution.

Choose one of these assumptions and think of a way that the researchers could — or did — build into the study a way to test whether the assumption seems valid.



- 7) Some people say that aquarium collecting is not a problem on Maui, while others believe that it is a problem in some areas or could quickly become one. Write one paragraph about what you would do to find out whether aquarium collecting is a threat to Maui reef animals. Write another paragraph about what you think should be done, if anything, to protect Maui reef fish populations from the impacts of collecting. (If you played the “Weren’t There More of Us?” game, how did what you learned from that game influence your response to these questions?)

